
Figure
Extracellular Stimuli
(Mitogeas, Stress Signels, etc.)
Cell Membrane
Cytoplan U
MAPKK activation
MAPK activation
Nucleus Pou
Active Transactivation Transactivation Transactivation Transactivation
Transcrip. Fector
·
— form
·

· ·	. <u>.</u>	To Tamba Ma	
		DBD: 72	<u>-</u>
	٠ كَا لِمَ اللَّهُ * فَيْ * مَا لَمُ اللَّهُ * فَيْ *		<u>.</u>
	7 × × × × × × × × × × × × × × × × × × ×	4 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	<u>.J</u>
	- S O O O O O O O O O O O O O O O O O O	-	
· .	\$\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	+	
	1 5 5		\$
		17	<u> </u>
	₹.		<u> </u>
	₹ <u>√</u>	C. rank	1,1
		<u> </u>	a—
		1 20	7-
		*	
		\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	
			_
		- Contract	
		+ %	
Conmon			
Lonnon			
			

4.1.1. pFR-Luc Plasmid



Sequence of GAL4 Binding Element in the pFR-Luc Plasmid

GT CGGAGTACTGTCCTCCG AG CGGAGTACTGTCCTCCG


AG CGGAGTACTGTCCTCCG AG CGGAGTACTGTCCTCCG

AG CGGAGTACTGTCCTCCG AG CGGAGACTCTAGAGGG

TATATATGGATCCCCGGGT AC CGAGCTCGAATTC--

-- CAGCTTGGCATTCCGGTACTGTTGGTAAMTG--Luciferase

4.1.2. Fusion Transactivator Plasmids



4.1.3. Control Plasmids






Figure 7 will tailor to each application

pFA-CMV Plasmid



4.2. Preparation of medium and reagents

Luciferase Assay Reagent (1×) 40.0 mM tricine (pH 7.8) 0.5 mM ATP 10 mM MgSO₄ 0.5 mM EDTA 10.0 mM DTT 0.5 mM coenzyme A 0.5 mM luciferin Cell Lysis Buffer (5×) 40 mM tricine (pH 7.8) 50 mM NaCl 2 mM EDTA 1 mM MgSO₄ 5 mM DTT 1% Triton® X-100






Figure 9




Figure 10